arXiv:1302.3695v1 [math.NT] 15 Feb 2013

ZEROS OF PARTIAL SUMS OF POWER SERIES FOR A FAMILY OF EXPONENTIAL INTEGRALS, I

ANTONIO R. VARGAS

Abstract. We are interested in studying the asymptotic behavior of the zeros of partial sums of power series for a family of entire functions defined by exponential integrals. The zeros grow on the order of $O(n)$, and after rescaling we explicitly calculate their limit curve. We find that the rate that the zeros approach the curve depends on the order of the singularities/zeros of the integrand in the exponential integrals. As an application of our findings we derive results concerning the zeros of partial sums of power series for Bessel functions of the first kind.

1. Introduction

In this paper we are concerned with the asymptotic behavior of the zeros of the polynomial sequence given by the partial sums of a convergent power series. If f is a function which is analytic at the origin, then it can be represented by a power series

$$f(z) = \sum_{k=0}^{\infty} a_k z^k$$

which converges near $z = 0$. We denote the n^{th} partial sum of this power series by

$$s_n[f](z) = \sum_{k=0}^{n} a_k z^k, \tag{1}$$

and we refer to $s_n[f](z)$ as the n^{th} section of f.

If the power series for f has a finite radius of convergence, then it is a classical result of Jentzsch [8] that every point on the circle of convergence of the power series will be a limit point of the zeros of the sections $s_n[f](z)$. In addition, because power series converge uniformly on compact subsets of their domains of convergence, Hurwitz's theorem (see, e.g., [9, p. 4]) tells us that any zero of f inside the radius of convergence will also be a limit point of the zeros of the sections.

The behavior of the zeros becomes much more interesting when f is entire. The topic can be traced back to Szegő, who studied in [14] the sections of the exponential function e^z, given by

$$s_n[\exp](z) = \sum_{k=0}^{n} \frac{z^k}{k!}.$$

Szegő found that the zeros of the normalized sections $s_n[\exp](nz)$ have as their set of limit points the simple closed loop

$$D = \left\{ z \in \mathbb{C} : |z| \leq 1 \text{ and } \left| z e^{1-z} \right| = 1 \right\}.$$

This is often referred to as the Szegő curve. The rate that the zeros approach this curve was first studied by Buckholtz [5], who showed that every zero of $s_n[\exp](nz)$ lies within a distance of $2e/\sqrt{n}$ of D. Carpenter, Varga, and Waldvogel [6] examined this phenomenon in detail and showed that Buckholtz's result gives the best-possible asymptotic order. The statement of the theorem involves the complementary error function

$$\operatorname{erfc} z = \frac{2}{\sqrt{\pi}} \int_z^\infty e^{-t^2}\, dt,$$

where the path of integration begins at z and travels to the right to ∞.

Theorem 1.1 (CVW). *For $\Omega \subseteq \mathbb{C}$, define* $\operatorname{maxdist}(\Omega, C) = \sup_{z\in\Omega}\{\operatorname{dist}(z, C)\}$. *If $\{z_{k,n}\}_{k=1}^n$ are the zeros of $s_n[\exp](nz)$ and if t_1 is the zero of the complementary error function* erfc *closest to the origin in the upper half-plane, then*

$$\liminf_{n\to\infty} \sqrt{n}\cdot \operatorname{maxdist}(\{z_{k,n}\}_{k=1}^n, D) \geq \Re(t_1) + \Im(t_1) \approx 0.636657.$$

The authors also showed that the zeros which are bounded away from the point $z = 1$ approach the Szegő curve more quickly.

Theorem 1.2 (CVW). *Let C_δ be the ball of radius δ centered at $z = 1$. If $\{z_{k,n}\}_{k=1}^n$ are the zeros of $s_n[\exp](nz)$ and if δ is any fixed number with $0 < \delta \leq 1$, then*

$$\operatorname{maxdist}(\{z_{k,n}\}_{k=1}^n \setminus C_\delta,\, D) = O\left(\frac{\log n}{n}\right)$$

as $n \to \infty$.

There have been some similar results for other power series, notably sine and cosine (see [15] and the references therein), a class of confluent hypergeometric functions [11], finite sums of exponentials [2], and even some classes of divergent power series [7]. For a survey of this topic the reader is referred to [16].

In this paper we will continue in this vein, studying power series defined by exponential integrals of a certain form, which we introduce in Section 2. Special cases of these functions include the aforementioned confluent hypergeometric functions as well as Bessel functions of the first kind, the latter of which we will discuss in detail in Section 4.

Our results were particularly inspired by the work of Norfolk [11] on the confluent hypergeometric functions, defined by

$${}_1F_1(1; b; z) = \Gamma(b) \sum_{k=0}^\infty \frac{z^k}{\Gamma(k+b)}.$$

Norfolk studied the case where b is real and $b \neq 1, 0, -1, -2, \ldots$ (the function is defined by analytic continuation when $b \leq 0$). His main tool was an exponential integral representation of the functions valid for $b > 1$ (see Section 4). The extension of some of Norfolk's results to the case where b is complex with $\Re(b) > 1$ will be a consequence of our main theorems.

This paper is the first in a series of two. The results within were originally obtained in the author's Master's thesis [16] and form an analogue to Theorem 1.2. The second paper in the series will contain new results; in particular an analogue to Theorem 1.1.

2. Definitions and preliminaries

The functions we are interested in are defined by integrals of the form $\int_{-a}^b \varphi(t) e^{zt}\, dt$. The restrictions we place on the function φ are determined essentially by the abilities of Watson's lemma, discussed below.

Suppose $0 \le a, b < \infty$ and let $\varphi : [-a, b] \to \mathbb{C} \cup \{\infty\}$ be a function satisfying

$$\int_{-a}^{b} |\varphi(t)|\, dt < \infty$$

and $\varphi(t) = (t+a)^{\mu} f_1(t+a) = (b-t)^{\nu} f_2(b-t)$, where

(1) $\mu, \nu \in \mathbb{C}$ with $\Re(\mu) > -1$ and $\Re(\nu) > -1$,
(2) $f_1, f_2 : [0, a+b] \to \mathbb{C} \cup \{\infty\}$ with $f_1(0)$ and $f_2(0)$ both finite and nonzero,
(3) in a neighborhood of $t = 0$, both $f_1'(t)$ and $f_2'(t)$ exist and are bounded.

Define

$$F(z) = \int_{-a}^{b} \varphi(t) e^{zt}\, dt.$$

It is a consequence of the dominated convergence theorem that the function F is entire, and its sections are given by the formula

$$s_n[F](z) = \sum_{k=0}^{n} \frac{z^k}{k!} \int_{-a}^{b} \varphi(t) t^k\, dt. \tag{2}$$

We can view $F(z)$ as the exponential generating function of these particular integral moments of φ.

Properties (1), (2), and (3) above describe how the function φ behaves near the endpoints of integration, and perhaps more importantly they name various quantities we will refer to throughout the paper. Property (3) serves a special purpose: it allows us to determine a simple error term in the asymptotic expansion of F through the use of Watson's lemma (Theorem 2.2).

Next we collect some theorems which will aid us in proving our results. The first such theorem is due to Rosenbloom [12, 13].

A sequence of sections $\{s_N[f](z)\}$ is said to have a positive fraction of zeros in any sector with vertex at the origin if

$$\liminf_{N\to\infty} \frac{\sharp_N^{\angle}(\theta_1, \theta_2)}{N} > 0$$

for any fixed θ_1 and θ_2, where $\sharp_N^{\angle}(\theta_1, \theta_2)$ is the number of zeros of the section $s_N[f](z)$ in the sector $\theta_1 \le \arg z \le \theta_2$. Rosenbloom's result is as follows.

Theorem 2.1 (Rosenbloom)**.** *Let*

$$f(z) = \sum_{k=0}^{\infty} a_k z^k$$

be an entire function of finite positive order ρ and let $s_n[f](z)$ be as in equation (1). Define $\sharp_n^{\circ}(R)$ to be the number of zeros of $s_n[f](z)$ in the disk $|z| \le R$ and define $\rho_n = |a_n|^{-1/n}$. For any increasing sequence of indices $\{N\}$ such that the sequence of sections $\{s_N[f](z)\}$ has a positive fraction of zeros in any sector with vertex at the origin, the existence of which is guaranteed, and for $0 \le r < 1$ and $\epsilon > 0$ we have

$$\liminf_{N\to\infty} \frac{\sharp_N^{\circ}\Big((e^{1/\rho} + \epsilon)\rho_N\Big) - \sharp_N^{\circ}(\rho_N r)}{N} \ge 1 - r^{\rho} > 0.$$

Further, the number of zeros of $s_N[f](z)$ satisfying

$$|\rho_N z| > e^{1/\rho} + \epsilon$$

is bounded.

We will also require Watson's lemma on the asymptotic behavior of exponential integrals. We refer the reader to [10] for a thorough discussion of this result. In the following, λ is a complex parameter.

Theorem 2.2 (Watson's lemma)**.** *Suppose* $0 < T \leq \infty$ *and* $\varphi : [0, T] \to \mathbb{C} \cup \{\infty\}$ *is a function satisfying*

$$\int_0^T |\varphi(t)|\, dt < \infty$$

and $\varphi(t) = t^\sigma h(t)$*, where* $\Re(\sigma) > -1$*,* $h(0) \neq 0$*, and* $h'(t)$ *exists and is bounded in a neighborhood of* $t = 0$*. Then the exponential integral*

$$\Phi(\lambda) = \int_0^T \varphi(t) e^{-\lambda t}\, dt$$

is finite for all $\Re(\lambda) > 0$*, and*

$$\Phi(\lambda) = \frac{h(0)\Gamma(\sigma+1)}{\lambda^{\sigma+1}} + O\big(\lambda^{-\sigma-2}\big)$$

as $\lambda \to \infty$ *with* $|\arg \lambda| \leq \theta$ *for any fixed* $0 \leq \theta < \pi/2$*.*

Though this form of Watson's lemma only gives an asymptotic for $\Phi(\lambda)$ as $\lambda \to \infty$ to the right, it can easily be extended to address the case when $\lambda \to \infty$ to the left if we assume that T is finite.

Corollary 2.3. *Suppose* $0 < T < \infty$ *and* $\varphi : [0, T] \to \mathbb{C} \cup \{\infty\}$ *is a function satisfying*

$$\int_0^T |\varphi(t)|\, dt < \infty$$

and $\varphi(t) = (T-t)^\sigma h(T-t)$*, where* $\Re(\sigma) > -1$*,* $h(0) \neq 0$*, and* $h'(t)$ *exists and is bounded in a neighborhood of* $t = 0$*. Then the exponential integral*

$$\Phi(\lambda) = \int_0^T \varphi(t) e^{\lambda t}\, dt$$

is finite for all $\Re(\lambda) > 0$*, and*

$$\Phi(\lambda) = \frac{h(0)\Gamma(\sigma+1)}{\lambda^{\sigma+1}}\, e^{T\lambda} + O\big(\lambda^{-\sigma-2} e^{T\lambda}\big)$$

as $\lambda \to \infty$ *with* $|\arg \lambda| \leq \theta$ *for any fixed* $0 \leq \theta < \pi/2$*.*

Proof. We have

$$\begin{aligned} \Phi(\lambda) e^{-T\lambda} &= \int_0^T \varphi(t) e^{-\lambda(T-t)}\, dt \\ &= \int_0^T \varphi(T-u) e^{-\lambda u}\, du \\ &= \int_0^T u^\sigma h(u) e^{-\lambda u}\, du \\ &= \frac{h(0)\Gamma(\sigma+1)}{\lambda^{\sigma+1}} + O\big(\lambda^{-\sigma-2}\big) \end{aligned}$$

as $\lambda \to \infty$ with $|\arg \lambda| \leq \theta$ for any fixed $0 \leq \theta < \pi/2$, by Watson's lemma. □

3. Main results

Recall from Section 2 that we are concerned with functions of the form

$$F(z) = \int_{-a}^{b} \varphi(t) e^{zt}\, dt$$

with φ satisfying some light requirements. The n^{th} section of F is the polynomial

$$s_n[F](z) = \sum_{k=0}^{n} \frac{z^k}{k!} \int_{-a}^{b} \varphi(t) t^k\, dt.$$

The statements and proofs of the main results of this section depend on the relative sizes of a and b and of $\Re(\mu)$ and $\Re(\nu)$. To this end, define

$$c = \max\{a, b\}$$

and

$$\xi = \begin{cases} \Re(\mu) & \text{if } a > b, \\ \Re(\nu) & \text{if } a < b, \\ \min\{\Re(\mu), \Re(\nu)\} & \text{if } a = b. \end{cases}$$

Lastly, let $\{N\}$ be an increasing subsequence of the indices $\{n\}$ for which the sequence of sections $\{s_N[F](z)\}$ has a positive fraction of zeros in any sector with vertex at the origin (see Section 2). If $a = b$ and $\Re(\mu) = \Re(\nu)$, we also impose the condition that the indices $\{N\}$ are chosen so that quantity

$$(-1)^N f_1(0)\Gamma(\mu+1) + f_2(0)\Gamma(\nu+1) a^{\nu-\mu} N^{\mu-\nu} \tag{3}$$

is bounded away from 0. Such a subsequence is guaranteed to exist by Theorem 2.1. This condition imposed on the quantity in (3) ensures that we can use the asymptotic representations derived in Lemma 3.5 and Lemma 3.6 without incident.

Theorem 3.1. *Let F be an exponential integral function as in Section 2 and let $\{N\}$ be a subsequence of the indices $\{n\}$ as defined above.*

- *(i) Let $\{z_N\}$ be a sequence of complex numbers such that $s_N[F](Nz_N) = 0$ for all N which has a limit point in the region $-1/a < Re(z) < 0$. Then the elements of the sequence satisfy*

  $$\left|c z_N e^{1+az_N}\right| = 1 + \left(\xi - \Re(\mu) + \frac{1}{2}\right)\frac{\log N}{N} + O(1/N)$$

  as $N \to \infty$.
- *(ii) Let $\{z_N\}$ be a sequence of complex numbers such that $s_N[F](Nz_N) = 0$ for all N which has a limit point in the region $0 < \Re(z) < 1/b$. Then the elements of the sequence satisfy*

  $$\left|c z_N e^{1-bz_N}\right| = 1 + \left(\xi - \Re(\nu) + \frac{1}{2}\right)\frac{\log N}{N} + O(1/N)$$

  as $N \to \infty$.

Theorem 3.2. *It is true that*

(i) Every point on the curve

$$D_{a,b} = \left\{ z \in \mathbb{C} : \Re(z) \le 0,\ \ |z| \le \frac{1}{c},\ \ \textit{and}\ \left| cze^{1+az} \right| = 1 \right\}$$
$$\cup \left\{ z \in \mathbb{C} : \Re(z) \ge 0,\ \ |z| \le \frac{1}{c},\ \ \textit{and}\ \left| cze^{1-bz} \right| = 1 \right\}$$

is a limit point of the zeros of the sections $s_N[F](Nz)$.

(ii) The limit points of the zeros of the sections $s_N[F](Nz)$ *on the imaginary axis are precisely the points on the line segment*

$$D_{\mathrm{imag}} = \left\{ z \in \mathbb{C} : \Re(z) = 0\ \ \textit{and}\ \ |z| \le \frac{1}{ec} \right\}.$$

(iii) All but finitely many limit points of the zeros of the sections $s_N[F](Nz)$ *lie on* $D_{a,b} \cup D_{imag}$*. Any limit points not on* $D_{a,b} \cup D_{imag}$ *lie outside of the circle* $|z| = 1/c$.

Remark 3.3. The expressions in parts (*i*) and (*ii*) of Theorem 3.1 give information about whether the zeros eventually lie on the inside or the outside of the limit curve based on signs of the quantities $\xi - \Re(\mu) + 1/2$ and $\xi - \Re(\nu) + 1/2$. For example, if $\xi - \Re(\mu) + 1/2 > 0$, then it will eventually be true that the zeros which approach $D_{a,b}$ in the left half-plane will satisfy $\left| cze^{1+az} \right| > 1$ and hence will lie outside of $D_{a,b}$. If either of the quantities $\xi - \Re(\mu) + 1/2$ or $\xi - \Re(\nu) + 1/2$ is zero then the theorem does not give any information about the direction from which the zeros approach the relevant part of the curve.

Figure 1 and Figure 2 illustrate two interesting cases of the results in Theorem 3.1 and Theorem 3.2. The reader may refer to [16] for more plots of this kind.



FIGURE 1. Zeros of the normalized section $s_{80}[F](80z)$ and the limit curve $D_{a,b} \cup D_{\mathrm{imag}}$ with $a = b = 1$ and $\varphi(t) = (1-t)^{3/2}(1+t)^{-1/2+i}$. Note that $\xi - \Re(\nu) + 1/2 = -3/2 < 0$, which predicts that the zeros in the right half-plane will approach the limit curve from the interior.

To prove these theorems we will require a few lemmas, the first of which concerns the asymptotic behavior of the function F.



FIGURE 2. Zeros of the normalized section $s_{80}[F](80z)$ and the limit curve $D_{a,b} \cup D_{\mathrm{imag}}$ with $a = 1$, $b = \frac{21}{20}$, and $\varphi(t) = (\frac{21}{20} - t)^i (1+t)^{1/2-i}$. Note that $\xi - \Re(\mu) + 1/2 = 0$, which predicts that the zeros in the left half-plane will approach the limit curve at a rate of $O(1/n)$.

Lemma 3.4. *As* $n \to \infty$,

$$F(nz) = f_1(0)\Gamma(\mu+1)(-nz)^{-\mu-1}e^{-anz}\Big(1 + O(1/n)\Big)$$

when z *is restricted to a compact subset of* $\Re(z) < 0$, *and*

$$F(nz) = f_2(0)\Gamma(\nu+1)(nz)^{-\nu-1}e^{bnz}\Big(1 + O(1/n)\Big)$$

when z *is restricted to a compact subset of* $\Re(z) > 0$.

Proof. This follows from a direct application of Corollary 2.3. To see this, suppose first that z is restricted to a compact subset of $\Re(z) < 0$, and make the substitution $t = b - s$ in the integral for $F(nz)$ to get

$$F(nz) = \int_{-a}^{b} \varphi(t)e^{nzt}\,dt = e^{bnz}\int_0^{a+b} \varphi(b-s)e^{-nzs}\,ds,$$

which, after replacing z with $-z$, is of the form required by the corollary. Next, suppose that z is restricted to a compact subset of $\Re(z) > 0$, and make the substitution $t = s - a$ in the definition of $F(nz)$ to get

$$F(nz) = \int_{-a}^{b} \varphi(t)e^{nzt}\,dt = e^{-anz}\int_0^{a+b} \varphi(s-a)e^{nzs}\,ds,$$

which is also of the required form. □

We must also find asymptotics for the integral moments of φ and hence for the power series coefficients of F.

Lemma 3.5. *We have*

$$\int_{-a}^{b} \varphi(t) t^n \, dt = (-1)^n f_1(0) \Gamma(\mu+1) n^{-\mu-1} a^{n+\mu+1} + O\big(n^{-\mu-2} a^n\big) \\ + f_2(0)\Gamma(\nu+1) n^{-\nu-1} b^{n+\nu+1} + O\big(n^{-\nu-2} b^n\big)\,.$$

as $n \to \infty$.

Proof. If $a \neq 0$ we calculate

$$\begin{aligned} \int_{-a}^{0} \varphi(t) t^n \, dt &= (-a)^n \int_{-a}^{0} \varphi(t) e^{n \log(-t/a)} \, dt \\ &= (-a)^n \int_{0}^{a} \varphi(s-a) e^{n \log(1-s/a)} \, ds \\ &= (-a)^n \int_{0}^{a} s^\mu f_1(s) e^{n \log(1-s/a)} \, ds. \end{aligned}$$

Letting $s = a(1-e^{-r})$ gives

$$\begin{aligned} \int_{-a}^{0} \varphi(t) t^n \, dt &= (-1)^n a^{n+\mu+1} \int_{0}^{\infty} (1-e^{-r})^\mu f_1(a - ae^{-r}) e^{-r} e^{-nr} \, dr \\ &= (-1)^n a^{n+\mu+1} \int_{0}^{\infty} r^\mu \psi_a(r) e^{-nr} \, dr, \end{aligned}$$

where

$$\psi_a(r) = \left(\frac{1-e^{-r}}{r}\right)^\mu f_1(a - ae^{-r}) e^{-r}$$

has a bounded derivative in a neighborhood of $r = 0$. We may now apply Watson's lemma to conclude that

$$\begin{aligned} \int_{-a}^{0} \varphi(t) t^n \, dt &= (-1)^n \psi_a(0) \Gamma(\mu+1) n^{-\mu-1} a^{n+\mu+1} + O\big(n^{-\mu-2} a^n\big) \\ &= (-1)^n f_1(0) \Gamma(\mu+1) n^{-\mu-1} a^{n+\mu+1} + O\big(n^{-\mu-2} a^n\big)\,. \end{aligned}$$

Using an identical argument we find that

$$\int_{0}^{b} \varphi(t) t^n \, dt = f_2(0) \Gamma(\nu+1) n^{-\nu-1} b^{n+\nu+1} + O\big(n^{-\nu-2} b^n\big)\,,$$

which completes the proof. □

A similar argument can be used to prove the following.

Lemma 3.6.

$$\int_{-a}^{b} \frac{\varphi(t)}{1-zt} t^{n+1} \, dt = (-1)^{n+1} \frac{f_1(0)\Gamma(\mu+1)}{1+az} n^{-\mu-1} a^{n+\mu+2} + O\big(n^{-\mu-2} a^n\big) \\ + \frac{f_2(0)\Gamma(\nu+1)}{1-bz} n^{-\nu-1} b^{n+\nu+2} + O\big(n^{-\nu-2} b^n\big)$$

as $n \to \infty$ *uniformly when* z *is restricted to a compact subset of the doubly-punctured plane* $\{z \in \mathbb{C} : z \neq -1/a \;\text{ and }\; z \neq 1/b\}$.

In the next lemma we will use the result of Lemma 3.5 to place a bound on where the limit points of the zeros of the normalized sections $s_N[F](Nz)$ may lie.

Lemma 3.7. *All but finitely many limit points of the zeros of the sections* $s_N[F](Nz)$ *lie in the disk* $|z| \le 1/c$*, where* $c = \max\{a, b\}$*.*

Proof. From Stirling's formula

$$n! \sim \left(\frac{n}{e}\right)^n \sqrt{2\pi n}$$

we have

$$(n!)^{1/n} \sim \frac{n}{e}$$

as $n \to \infty$, and with the aid of Lemma 3.5 we calculate

$$\left|\int_{-a}^{b} \varphi(t) t^N \, dt\right|^{-1/N} \longrightarrow \frac{1}{c}$$

as $N \to \infty$, where the subsequence of indices $\{N\}$ is as defined above Theorem 3.1. Combining these we see that the power series coefficients of $F(z)$, which are given by

$$a_n = \frac{1}{n!} \int_{-a}^{b} \varphi(t) t^n \, dt,$$

satisfy

$$\rho_N = |a_N|^{-1/N} \sim \frac{N}{ec},$$

where ρ_N is as defined in Theorem 2.1. Thus the order ρ of F is calculated to be

$$\rho = \limsup_{n\to\infty} \frac{\log n}{\log \rho_n} = 1,$$

where all indices n are taken into account (see, e.g., [3, p. 9]).

By Theorem 2.1 we have, for any $\epsilon > 0$,

$$\liminf_{N\to\infty} \frac{\sharp_N^\circ\Big((1+\epsilon)N/c\Big)}{N} = 1,$$

where $\sharp_N^\circ\big((1+\epsilon)N/c\big)$ is the number of zeros of $s_N[F](z)$ in the disk $|z| \le (1+\epsilon)N/c$. From this we conclude that all but finitely many limit points of the zeros of the sections $s_N[F](Nz)$ lie in the disk $|z| \le 1/c$, as desired. □

We may now prove the first theorem in this section.

Proof of Theorem 3.1. By definition we have

$$F(nz) = \int_{-a}^{b} \varphi(t) e^{nzt} \, dt,$$

and

$$s_n[F](nz) = \int_{-a}^{b} \varphi(t) s_n[\exp](nzt) \, dt.$$

Subtracting these we get

$$\begin{aligned} F(nz) - s_n[F](nz) &= \int_{-a}^{b} \varphi(t) \left(e^{nzt} - s_n[\exp](nzt)\right) \, dt \\ &= \int_{-a}^{b} \varphi(t) e^{nzt} g_n(zt) \, dt, \qquad (4) \end{aligned}$$

where

$$g_n(z) = 1 - e^{-nz} s_n[\exp](nz).$$

It was shown by Szegő in [14] (see also [6], [4], and [11]) that

$$g_n(z) = \frac{\left(ze^{1-z}\right)^n}{\sqrt{2\pi n}} \cdot \frac{z}{1-z}\Big(1-\epsilon_n(z)\Big), \tag{5}$$

where $\epsilon_n(z) = O(1/n)$ as $n \to \infty$ uniformly when z is restricted to a compact subset of $\Re(z) < 1$. Upon substituting this into equation (4) we get

$$F(nz) - s_n[F](nz) = \frac{e^n z^{n+1}}{\sqrt{2\pi n}} \int_{-a}^{b} \frac{\varphi(t)}{1-zt}\, t^{n+1}\Big(1-\epsilon_n(zt)\Big)\, dt.$$

If $c = \max\{a, b\}$ it follows that zeros of $s_n[F](nz)$ which remain in compact subsets of the region $|\Re(z)| < 1/c$ satisfy

$$F(nz) = \frac{e^n z^{n+1}}{\sqrt{2\pi n}} \int_{-a}^{b} \frac{\varphi(t)}{1-zt}\, t^{n+1}\, dt\, \Big(1 + O(1/n)\Big) \tag{6}$$

as $n \to \infty$, where we have used Lemma 3.6 to bring the error term outside of the integral.

Let $\{N\}$ be a sequence of indices as defined above Theorem 3.1. We showed in Lemma 3.7 that all but finitely many the limit points of the zeros of the sections $s_N[F](Nz)$ lie in the disk $|z| \le 1/c$. Define

$$\Delta = \left\{ z \in \mathbb{C} : |z| \le \frac{1}{c}, \quad z \ne -\frac{1}{a}, \quad \text{and} \quad z \ne \frac{1}{b} \right\}.$$

Suppose first that $\{z_N\}$ is a sequence in $\mathbb{C}$ such that $s_N[F](Nz_N) = 0$ for all N and such that the sequence has a limit point in $\Delta \cap \{z \in \mathbb{C} : \Re(z) < 0\}$. Note that this implies there is a $\delta > 0$ such that $|z_N + 1/a| > \delta$ for N large enough.

It follows from Lemma 3.4 that

$$|F(Nz_N)|^{1/N} = |e^{-az_N}| \left(1 - (\Re(\mu)+1)\frac{\log N}{N} + O(1/N)\right) \tag{7}$$

as $N \to \infty$, and if

$$\xi = \begin{cases} \Re(\mu) & \text{if } a > b, \\ \Re(\nu) & \text{if } a < b, \\ \min\{\Re(\mu), \Re(\nu)\} & \text{if } a = b, \end{cases}$$

then we have from Lemma 3.6 that

$$\left| \frac{e^N z^{N+1}}{\sqrt{2\pi N}} \int_{-a}^{b} \frac{\varphi(t)}{1-zt}\, t^{N+1}\, dt \right|^{1/N} = |ecz_N| \left(1 - \left(\xi + \frac{3}{2}\right)\frac{\log N}{N} + O(1/N)\right) \tag{8}$$

as $N \to \infty$. Upon substituting equations (7) and (8) into equation (6) we see that these zeros z_N satisfy

$$\left|cz_N e^{1+az_N}\right| = 1 + \left(\xi - \Re(\mu) + \frac{1}{2}\right)\frac{\log N}{N} + O(1/N)$$

as $N \to \infty$, which proves part (i) of Theorem 3.1.

Suppose now that $\{z_N\}$ is a sequence such that $s_N[F](Nz_N) = 0$ for all N and such that the sequence has a limit point in $\Delta \cap \{z \in \mathbb{C} : \Re(z) > 0\}$. Note that this implies there is a $\delta > 0$ such that $|z_N - 1/b| > \delta$ for N large enough.

Here it follows from Lemma 3.4 that

$$|F(Nz_N)|^{1/N} = |e^{bz_N}| \left(1 - (\Re(\nu)+1)\frac{\log N}{N} + O(1/N)\right)$$

as $N \to \infty$. Subtituting this and equation (8) into equation (6) we see that these zeros z_N satisfy

$$\left| c z_N e^{1-b z_N} \right| = 1 + \left(\xi - \Re(\nu) + \frac{1}{2} \right) \frac{\log N}{N} + O(1/N)$$

as $N \to \infty$, which proves part (ii) of Theorem 3.1. □

We will now use this result to prove the second theorem in this section.

Proof of Theorem 3.2. It follows from Lemma 3.7 and Theorem 3.1 that all but finitely many limit points of the zeros of the sections $s_N[F](Nz)$ with $\Re(z) \neq 0$ must lie on the curve $D_{a,b}$ as defined in part (i) of Theorem 3.2. That every point on $D_{a,b}$ is such a limit point follows from the choice of the subsequence $\{N\}$ which ensures that the sequence of sections $\{s_N[F](z)\}$ has a positive fraction of zeros in any sector with vertex at the origin. It is straightforward to show that for any $0 \leq \theta < 2\pi$ there is a unique $r > 0$ such that $re^{i\theta} \in D_{a,b}$. This proves part (i) of Theorem 3.2.

From the asymptotic expansion for $F(nz)$ in Lemma 3.4 we see that the limit points of the zeros of $F(nz)$ must lie on the imaginary axis. Further, we claim that F has infinitely many zeros, so every point on the imaginary axis is such a limit point. Indeed, if F had finitely many zeros then, since F has order 1, the Hadamard factorization theorem would tell us that we can find a constant a and a polynomial p such that

$$F(z) = e^{az} p(z). \tag{9}$$

But if $y \in \mathbb{R}$ then

$$|F(iy)| = \left| \int_{-a}^{b} \varphi(t) e^{iyt}\, dt \right| \leq \int_{-a}^{b} |\varphi(t)|\, dt.$$

Taking the limit as $z \to i\infty$ in (9) we see that we must have $\Im(a) > 0$, and taking the limit as $z \to -i\infty$ we must have $\Im(a) < 0$, a contradiction. Thus the claim is verified.

If $\{z_N\}$ is a sequence of complex numbers such that $s_N[F](Nz_N) = 0$ for all N which has a limit point on $\{z \in \mathbb{C} : \Re(z) = 0 \text{ and } \Im(z) > 1/(ec)\}$ then by equation (6) and Lemma 3.6 we must have $F(Nz_N) \to \infty$. We just showed that $F(iny)$ is bounded, so such a sequence of zeros cannot exist. Hence any limit points on the imaginary axis must satisfy $\Im(z) \leq 1/(ec)$.

Further, if $\{z_N\}$ is a sequence of zeros which has a limit point in $|z| \leq 1/(ec)$, then by equation (6) and Lemma 3.6 we must have $F(Nz_N) \to 0$. In other words, the zeros $\{z_N\}$ must approximate the zeros of $F(Nz)$. Thus part (ii) is proved.

The truth of part (iii) follows from combining the facts that a) all but finitely many limit points with $\Re(z) \neq 0$ must lie on $D_{a,b}$ and b) any limit points with $\Re(z) = 0$ must lie on D_{imag}. This completes the proof of Theorem 3.2. □

4. SPECIAL CASES OF THE EXPONENTIAL INTEGRALS

We mentioned in the introduction that the confluent hypergeometric functions with $b > 1$ studied by Norfolk [11] are a special case of the exponential integrals studied in this paper. Indeed, when $b > 1$ we have the integral represenation

$${}_1F_1(1; b; z) = (b-1) \int_0^1 (1-t)^{b-2} e^{zt}\, dt.$$

Our result extends some of Norfolk's results to the case of complex b with $\Re(b) > 1$.

Another important special case of the exponential integrals is the class of Bessel functions of the first kind, defined by

$$J_\alpha(z) = \left(\frac{z}{2}\right)^\alpha \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma(k+1)\Gamma(k+\alpha+1)} \left(\frac{z}{2}\right)^{2k}.$$

For $\Re(\alpha) > -1/2$ we have Poisson's integral representation (see, e.g., [17])

$$J_\alpha(z) = \frac{\left(\frac{z}{2}\right)^\alpha}{\Gamma\left(\alpha+\frac{1}{2}\right)\Gamma\left(\frac{1}{2}\right)} \int_{-1}^{1} \left(1-t^2\right)^{\alpha-1/2} e^{izt}\,dt,$$

which is of the required form once z is replaced with $-iz$. As far as we can tell, asymptotics for the zeros of sections of the Bessel functions have not been previously studied. We state the result formally as a corollary to Theorem 3.1 and Theorem 3.2. This corollary is illustrated in Figure 3.

Corollary 4.1. *Let J_α be the Bessel function of order α with $\Re(\alpha) > -1/2$, and for n even let*

$$s_n[J_\alpha](z) = \left(\frac{z}{2}\right)^\alpha \sum_{k=0}^{n/2} \frac{(-1)^k}{\Gamma(k+1)\Gamma(k+\alpha+1)} \left(\frac{z}{2}\right)^{2k}$$

be its n^{th} section. Then, for a subsequence $\{N\}$ of the indices $\{n\}$ as defined in Theorem 2.1, all but finitely many limit points of the zeros of the normalized sections $s_N[J_\alpha](Nz)$ lie on

$$\begin{aligned} D(J) = &\left\{z \in \mathbb{C} : \Im(z) \geq 0,\ |z| \leq 1,\ \textit{and}\ \left|ze^{1+iz}\right| = 1\right\} \\ &\cup \left\{z \in \mathbb{C} : \Im(z) \leq 0,\ |z| \leq 1,\ \textit{and}\ \left|ze^{1-iz}\right| = 1\right\} \\ &\cup \left\{x \in \mathbb{R} :\ -1/e \leq x \leq 1/e\right\}. \end{aligned}$$

Conversely, every point of $D(J)$ is a limit point of zeros. Zeros of $s_N[J_\alpha](Nz)$ in $\Im(z) > 0$ which do not converge to the point $z = i$ satisfy

$$\left|ze^{1+iz}\right| = 1 + \frac{\log N}{2N} + O(1/N)$$

as $N \to \infty$, and zeros in $\Im(z) < 0$ which do not converge to $z = -i$ satisfy

$$\left|ze^{1-iz}\right| = 1 + \frac{\log N}{2N} + O(1/N)$$

as $N \to \infty$. Furthermore, if α is real, the modified indices $\{N\}$ in this result may be replaced everywhere by the original even indices $\{n\}$ and, in this case, all limit points of the zeros of the normalized sections $s_n[J_\alpha](nz)$ lie on $D(J)$.

Proof. The only part of Corollary 4.1 which does not follow immediately from Theorem 3.1 or Theorem 3.2 is the claim that if α is real, the modified indices $\{N\}$ may be replaced by the original even indices $\{n\}$ and, in this case, all limit points of the zeros of the normalized sections $s_n[J_\alpha](nz)$ lie on $D(J)$.

To prove this claim we write

$$s_n[J_\alpha](-inz) = \left(-\frac{inz}{2}\right)^\alpha P_n\left(z^2\right),$$

where

$$P_n(z) = \sum_{k=0}^{n/2} \frac{n^{2k}}{4^k \Gamma(k+1)\Gamma(k+\alpha+1)} z^k.$$



FIGURE 3. LEFT: Zeros of the normalized section $s_{80}[J_0](80z)$ and the limit curve $D(J)$ as defined in Corollary 4.1.

By the Eneström-Kakeya theorem (see, e.g., [9, ch. 7] and [1]), all zeros of $P_n(z)$ satisfy

$$|z| \leq \frac{2n+4}{n^2} \cdot \frac{\Gamma\left(\frac{n}{2}+\alpha+2\right)}{\Gamma\left(\frac{n}{2}+\alpha+1\right)}.$$

We apply Stirling's formula for the Gamma function to see that

$$\frac{\Gamma\left(\frac{n}{2}+\alpha+2\right)}{\Gamma\left(\frac{n}{2}+\alpha+1\right)} \sim \frac{n}{2},$$

whence

$$\frac{2n+4}{n^2} \cdot \frac{\Gamma\left(\frac{n}{2}+\alpha+2\right)}{\Gamma\left(\frac{n}{2}+\alpha+1\right)} \longrightarrow 1$$

as $n \to \infty$. This tells us that the limit points of the zeros of the polynomials P_n, and hence of the sections $s_n[J_\alpha](nz)$, lie in the closed unit disk. This replaces the use of Theorem 2.1 and hence Lemma 3.7 in the proofs of Theorem 3.1 and Theorem 3.2. Indeed, Theorem 2.1 is the origin of the restriction of the indices to the subsequences $\{N\}$. This completes the proof of the corollary. $\square$

5. Discussion and future work

The exponential integrals in this paper were studied in part because of their simple definitions. It should be noted that their integral form facilitated symbolic manipulations and allowed for the use of standard asymptotic techniques. But they were also studied because their definition was flexible enough to allow the zeros of their sections to display some unique behavior, perhaps the most notable of which is the tendency for the zeros to approach the limit curve from the interior or the exterior depending on the orders of the singularities/zeros of the integrand φ at the endpoints of integration (see Remark 3.3).

As mentioned in Section 1, this paper is the first in a series of two. In the forthcoming second paper we will complement Theorem 3.1 by studying the asymptotics of the zeros which approach the points of the limit curve $D_{a,b}$ on the circle $|z| = 1/c$, which occur at $z = 1/b$ or $z = -1/a$ (or both). In general, the zeros approach these points more slowly than they approach the body of the curve. In fact, the unique behavior of the zeros described in the previous paragraph will play an important role in the analysis. Just as in this paper, results about the zeros of sections of the Bessel functions will appear as special cases.

References

1. N. Anderson, E. B. Saff, and R. S. Varga, *On the Eneström-Kakeya theorem and its sharpness*, Linear Algebra Appl. **28** (1979), 5–16.
2. P. Bleher and R. Mallison, Jr., *Zeros of sections of exponential sums*, Int. Math. Res. Not. (2006), Art. ID 38937, 49.
3. R. P. Boas, Jr., *Entire Functions*, Academic Press Inc., New York, 1954.
4. R. Boyer and W. M. Y. Goh, *On the zero attractor of the Euler polynomials*, Adv. in Appl. Math. **38** (2007), no. 1, 97–132.
5. J. D. Buckholtz, *A characterization of the exponential series*, Amer. Math. Monthly **73** (1966), no. 4, part II, 121–123.
6. A. J. Carpenter, R. S. Varga, and J. Waldvogel, *Asymptotics for the zeros of the partial sums of e^z. I*, Rocky Mountain J. Math. **21** (1991), no. 1, 99–120.
7. K. Dilcher and L. A. Rubel, *Zeros of sections of divergent power series*, J. Math. Anal. Appl. **198** (1996), no. 1, 98–110.
8. R. Jentzsch, *Untersuchungen zur Theorie der Folgen analytischer Funktionen*, Acta Math. **41** (1916), no. 1, 219–251.
9. M. Marden, *Geometry of Polynomials*, Second edition. Mathematical Surveys, No. 3, American Mathematical Society, Providence, R.I., 1966.
10. P. D. Miller, *Applied Asymptotic Analysis*, Graduate Studies in Mathematics, vol. 75, American Mathematical Society, Providence, RI, 2006.
11. T. S. Norfolk, *On the zeros of the partial sums to ${}_1F_1(1; b; z)$*, J. Math. Anal. Appl. **218** (1998), no. 2, 421–438.
12. P. C. Rosenbloom, *On sequences of polynomials, especially sections of power series*, Ph.D. thesis, Stanford University, 1944, Abstracts in Bull. Amer. Math. Soc. **48** (1942), 839; **49** (1943), 689.
13. ———, *Distribution of zeros of polynomials*, Lectures on Functions of a Complex Variable (W. Kaplan, ed.), The University of Michigan Press, Ann Arbor, 1955, pp. 265–285.
14. G. Szegő, *Über eine Eigenschaft der Exponentialreihe*, Berlin Math. Ges. Sitzungsber. **23** (1924), 50–64.
15. R. S. Varga and A. J. Carpenter, *Zeros of the partial sums of* $\cos(z)$ *and* $\sin(z)$*. III*, Appl. Numer. Math. **60** (2010), no. 4, 298–313.
16. A. R. Vargas, *Zeros of Sections of Some Power Series*, Master's thesis, Dalhousie University, 2012, arXiv:1208.5186 [math.NT].
17. G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1995, Reprint of the second (1944) edition.

Dept. of Mathematics and Statistics, Dalhousie University, Halifax, Nova Scotia B3H 4J5, Canada